(FILE 'HOME' ENTERED AT 14:05:03 ON 23 JUN 2004)

INDEX 'ADISCTI, ADISINSIGHT, ADISNEWS, AGRICOLA, ANABSTR, AQUASCI, BIOBUSINESS, BIOCOMMERCE, BIOSIS, BIOTECHABS, BIOTECHDS, BIOTECHNO, CABA, CANCERLIT, CAPLUS, CEABA-VTB, CEN, CIN, CONFSCI, CROPB, CROPU, DISSABS, DDFB, DDFU, DGENE, DRUGB, DRUGMONOG2, ...' ENTERED AT 14:07:02 ON 23 JUN 2004

SEA GLUCOAMYLASE

- 1 FILE ADISCTI
- 2 FILE ADISINSIGHT
- 700 FILE AGRICOLA
- 41 FILE ANABSTR
- 10 FILE AQUASCI
- 526 FILE BIOBUSINESS
- 48 FILE BIOCOMMERCE
- 2296 FILE BIOSIS
- 2085 FILE BIOTECHABS
- 2085 FILE BIOTECHDS
- 709 FILE BIOTECHNO
- 618 FILE CABA
- 29 FILE CANCERLIT
- 5385 FILE CAPLUS
- 633 FILE CEABA-VTB
 - 3 FILE CEN
- 29 FILE CIN
- 62 FILE CONFSCI
- 1 FILE CROPB
- 96 FILE DISSABS
- 22 FILE DDFB
- 23 FILE DDFU
- 1270 FILE DGENE
 - 22 FILE DRUGB
 - 39 FILE DRUGU
- 12 FILE EMBAL 1041 FILE EMBASE
- 562 FILE ESBIOBASE
 - OZ FILE ESBIODA
 - 9 FILE FEDRIP 2 FILE FOMAD
- 32 FILE FOREGE
- 461 FILE FROSTI
- 1396 FILE FSTA
- 549 FILE GENBANK
 - 2 FILE HEALSAFE
- 540 FILE IFIPAT
- 802 FILE JICST-EPLUS
- 937 FILE LIFESCI
- 1007 FILE MEDLINE
 - 2 FILE NIOSHTIC
 - 29 FILE NTIS
 - 3 FILE OCEAN
- 1347 FILE PASCAL
 - 2 FILE PHAR
 - 2 FILE PHIN
 - 68 FILE PROMT
 - 2 FILE PROUSDDR
- 1 FILE RDISCLOSURE
- 2119 FILE SCISEARCH 481 FILE TOXCENTER
- 5199 FILE USPATFULL
- 136 FILE USPAT2

1 FILE VETU

921 FILE WPIDS

L1

L2

2 FILE WPIFV

921 FILE WPINDEX

QUE GLUCOAMYLASE

FILE 'CAPLUS, BIOSIS, SCISEARCH, BIOTECHDS, FSTA, PASCAL, EMBASE, MEDLINE, LIFESCI, WPIDS, JICST-EPLUS, BIOTECHNO, AGRICOLA' ENTERED AT 14:08:08 ON 23 JUN 2004

1 S L1 AND (POSITION 402 OR AMINO ACID 402 OR TYR402 OR Y402)

ANSWER 143 OF 143 BIOTECHDS COPYRIGHT 2004 THOMSON DERWENT/ISI on STN 1.6 ACCESSION NUMBER: 1991-01384 BIOTECHDS

Expression and routeing of human lysosomal alpha-glucosidase

in transiently transfected mammalian cells;

glucoamylase gene cloning and expression in

COS-1 and HeLa cell culture; potential application in

glycogenosis type-II gene therapy

AUTHOR:

Hoefsloot L H; Willemsen R; Kroos M A; Hoogeveen-Westerveld

M; Hermans M M P; *Reuser A J J

LOCATION:

Department of Cell Biology and Genetics, Erasmus University,

P.O. Box 1738, 3000 DR, Rotterdam, The Netherlands.

SOURCE:

Biochem.J.; (1990) 272, 2, 485-92 CODEN: BIJOAK

DOCUMENT TYPE:

Journal

LANGUAGE:

English

cDNA encoding human acid alpha-glucosidase (glucoamylase, AB EC-3.2.1.3) was cloned in vector plasmid pSG5 and used to transform COS-1 and HeLa cells. mRNA was synthesized from the phage T7 promoter in front of the cDNA insert, and used to direct protein synthesis in a reticulocyte translation system. Only translation of sense mRNA led to protein production. Plasmid pSHAG1 did not encode a functional enzyme, due to an Arg residue replacing a Trp residue at position 402. The mutation did not affect enzyme production, but interfered with post-translational modification and intracellular transport of the precursor. Pulse-chase experiments suggested that the precursor was denatured. A Trp-402-containing enzyme (encoded by plasmid pSHAG2) was processed properly, was active, and reached the membrane and the medium. The proteins formed in the absence and in the presence of microsomes corresponded in their mol.weight to previously identified unglycosylated and glycosylated precursors of glucoamylase, obtained by translation in vitro of total RNA from human fibroblasts.

The enzyme may be useful in gene therapy of e.g. glycogenosis type II. (36 ref)

=> d 12 ibib ab

L2 ANSWER 1 OF 1 BIOTECHDS COPYRIGHT 2004 THOMSON DERWENT/ISI on STN

ACCESSION NUMBER: 1991-01384 BIOTECHDS

TITLE:

Expression and routeing of human lysosomal alpha-glucosidase

in transiently transfected mammalian cells;

glucoamylase gene cloning and expression in

COS-1 and HeLa cell culture; potential application in

glycogenosis type-II gene therapy

AUTHOR:

Hoefsloot L H; Willemsen R; Kroos M A; Hoogeveen-Westerveld

M; Hermans M M P; *Reuser A J J

LOCATION:

Department of Cell Biology and Genetics, Erasmus University,

P.O. Box 1738, 3000 DR, Rotterdam, The Netherlands.

SOURCE:

Biochem.J.; (1990) 272, 2, 485-92

CODEN: BIJOAK

DOCUMENT TYPE:

Journal English

LANGUAGE:

(36 ref)

AB cDNA encoding human acid alpha-glucosidase (glucoamylase,

EC-3.2.1.3) was cloned in vector plasmid pSG5 and used to transform COS-1 and HeLa cells. mRNA was synthesized from the phage T7 promoter in front of the cDNA insert, and used to direct protein synthesis in a reticulocyte translation system. Only translation of sense mRNA led to protein production. Plasmid pSHAG1 did not encode a functional enzyme, due to an Arg residue replacing a Trp residue at position 402. The mutation did not affect enzyme production, but interfered with post-translational modification and intracellular transport of the precursor. Pulse-chase experiments suggested that the precursor was denatured. A Trp-402-containing enzyme (encoded by plasmid pSHAG2) was processed properly, was active, and reached the membrane and the medium. The proteins formed in the absence and in the presence of microsomes corresponded in their mol.weight to previously identified unglycosylated and glycosylated precursors of glucoamylase, obtained by translation in vitro of total RNA from human fibroblasts. The enzyme may be useful in gene therapy of e.g. glycogenosis type II.







PMC Journals Book PubMed Nucleotide Protein Genome Structure **OMIM** Entrez Search PubMed Go Clear for ✓ Limits Preview/Index **Details** History Clipboard About Entrez Display. Abstract Sort Send to Text

Text Version

□ 1: Biochem J. 1990 Dec 1;272(2):485-92.

Related Articles, L

Entrez PubMed Overview Help | FAQ Tutorial New/Noteworthy E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
LinkOut
Cubby

Related Resources
Order Documents
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

Expression and routeing of human lysosomal alpha-glucosidase in transiently transfected mammalian cells.

Hoefsloot LH, Willemsen R, Kroos MA, Hoogeveen-Westerveld M, Hermans MM, Van der Ploeg AT, Oostra BA, Reuser AJ.

MGC-Department of Cell Biology and Genetics, Erasmus University, Rotterdam, The Netherlands.

Previously isolated lysosomal alpha-glucosidase cDNA clones were ligated to full-length constructs for expression in vitro and in mammalian cells. One of these constructs (pSHAG1) did not code for functional enzyme, due to an arginine residue instead of a tryptophan residue at amino acid position 402. The mutation does not affect the rate of enzyme synthesis, but interferes with post-translational modification and intracellular transport of the acid alpha-glucosidase precursor. Using immunocytochemistry it is demonstrated that the mutant precursor traverses the endoplasmic reticulum and the Golg complex, but does not reach the lysosomes. Pulse-chase experiments suggest premature degradation. The Trp-402-containing enzyme (encoded by construct pSHAG2) is proces properly, and has catalytic activity. A fraction of the enzyme is localized at the plasma membrane. It is hypothesized that membrane association of the acid alpha-glucosidase precursor, as demonstrated by Triton X-114 phase separation, is responsible for transpor this location. Transiently expressed acid alpha-glucosidase also enters the secretory pathway, since a catalytically active precursor is found in the culture medium. This precursor has the appropriate characteristics for use in enzyme replacement therapy. Efficient uptake via the mannose 6-phosphate receptor results in degradation of lysosom glycogen in cultured fibroblasts and muscle cells from patients with glycogenosis type II

PMID: 2268275 [PubMed - indexed for MEDLINE]

Display Abstract	Show: 20	Sort	Send to Text	Y
	Silow.			_

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer